PHYSICS-Part A

1.	If the them	distance between two masses	is doubled, the	gravitational attraction between
	(A)	Is doubled	(B)	Becomes four times
	(0)	Is reduced to half	(D)	Is reduced to a quarter
2.	If the	Kinetic energy of a particle in	linear motion i	s doubled, then its momentum will
	(A)	Remain unchanged	(B)	Increase $\sqrt{2}$ times
	(C)	Be quadrupled	(D)	Be doubled
3.	Rate	of doing work is called		
	(A)	Force	(B)	Acceleration
	(C)	Power	(D)	Displacement
3,	SI un	it of angular momentum is		
	(A)	rad/s ²	(B)	kg.m².s
	10)	- kg.m²/s	(D)	rad/s
5.	A co	oncave lens is kept in contact bination acts as a convex focal	with a conver length of 50 cm	k lens of focal length 20 cm. The h. The power of concave lens is
	(A)	-3D	(B)	+3D
	(C)	5D	(D)	6D
6.	Wha	t is the critical angle for a mate	erial for refracti	ve index $\sqrt{2}$.
	(A)	30°	(B)	45°
	(C)	60°	(D)	90°
7.	Foca	l length (F) and power (P) of p	olane glass plate	is
	-(A)	F=0, P=∞	(B)	F=∞, P=0
	(C)	4 F=1, P=1	(D)	F=0, P=0
8.	(A) (B) (C)	ch color deviates (i) most (ii) l Most for red and least for v Most for violet and least for same for both None for above	iolet	through a prism
	(D)			

9.	Ratio	of slit width, when amplitude o	f light waves e	manates from them have a ratio of			
	√7:√3. is						
	(A)	3:7	(B)				
	(C)	7:3	(D)	7: √3			
10.	Wher by	a wave undergoes a reflection	from rarer to d	enser medium, the phase changes			
	(A)	0°	(B)	π			
	(C)	π/2	(D)	-π			
11.	Dista (take	ance covered by a missile fired v g=10m/s ²) is	with initial velo	ocity of 300 m/s at an angle of 45°			
	(A)	90 km	(B)	9 km			
	(C)	81 km	(D)	30 km			
12.	Diele	ectric constant of a medium is 1	0. Its permittiv	ity will be			
	(A)	8.854×10 ⁻¹² C ² N ⁻¹ m ⁻²	(B)	$88.54 \times 10^{-12} \text{C}^2 \text{N}^{-1} \text{m}^{-2}$			
	(C)	0.8854×10 ⁻¹² C ² N ⁻¹ m ⁻²	(D)	None of the above			
13.	The	Kinetic energy and potential end in displacement is (amplitude = a	ergy of a partic	le executing S.H.M. will be equal,			
	(A)	a ² /2	(B)	$a/\sqrt{2}$			
	(C)	$a^2/\sqrt{2}$		a ² / 4			
14.	A sin	t t t t t t t t t t t t t t t t t t t	a different plan i.e $g_p = (1/100)$	net with its acceleration due to gravity g_E , then the time period of pendulum			
	(A)	100 times	(b)	10 times			
	(C)	1/10 times	(d)	1/100 times			
15.	The t in :	equation of wave is $y = 2 \sin \pi$ sec. The wave velocity is	(0.5x -200t) w	where x and y are expressed in cm and			
	(A)	100 cm/sec	(B)	200 cm/sec			
	-01	200 em/sec	(D)	400 cm/sec			
	Wor	done in moving the test charg	e from one po	int of equipotential surface to other is			
16.		infinite (∞)	(B)	zero			
	(A)		(D)	None of these			
	(C)	Unity					

		maxim	um penetrating po	wer:
17.	W	nich of following has maxim	(B)	Microwaves
	(A)	UV radiation	(D)	Radio waves
	(9)	Gamma rays	ross a color-coded	carbon resistor with first, second an rrent flowing through the resistor is
18.	AV	oltage of 200V is applied ac	low colors. The cu	
		10-14	(B)	3.33×10 ⁻⁴ A
	(A)	33.3×10 ⁻⁴ A	(D)	333×10 ⁻⁴ A
	(C)	33.3^10 A	$e x = a \sin (\omega t - \alpha t)$) and $y = b \cos (\omega t - \alpha)$. The phas
9.	Two	erene between the two is		
		0°	(B)	α°
	(A) (C)	90°	(D)	180°
).	circu	satellites S1 and S2 following lar orbits in same sense. The ectively. If radius of S1 is 10	heir periods of rev	revolving around a planet in coplanary volutions are one hour and 8 hours as of second satellite is
	(A)	4 × 10 ⁴ km	(B)	$0.4 \times 10^4 \text{ km}$
	(C)	40 × 10 ⁴ km	(D)	4 × 10 ⁴ m
		PHY	SICS - Part-	В
		(Attemp	t any 30 Questi	ons)
(Out of	speed (c), frequency (v) a	and wavelength (λ) on reflection
(A)	Speed varies while frequen	ncy and wavelengtl	remain same
0	B)	frequency and wavelength	vary while Speed	remains same
(C)	frequency varies while Spo	eed and wavelength	remain same
(1)	All remain same		
h		ple harmonic motion, which		
(1	4)	Kinetic energy is maximum	at mean position :	and minimum at extreme positions
	3)	Potential energy is maximum	m at extreme position	ons and minimum at mean position
	-)	Total energy always remain	s constant.	
1)))	All of the above		

21.

(A)	ρR	(B)	$R\sqrt{\rho}$
(C)	$\sqrt{\rho/R}$	(D)	1/pR
The f	orce between two free ele	ectrons spaced 1Å ap	art is
(A)	2.3×10 * N	(B)	0.23×10 ⁻⁸ N
(C)	4.6×10 ⁻⁸ N	(D)	230×10 ⁻⁸ N
			field $\vec{E} = 4\hat{i} + 4\hat{j} + 4\hat{k}$. The electric
(A)	40 units	(B)	200 units
(C)	20 units	(D)	400 units
(A)	40 V	(B)	200 V
(C)	15 V	(D)	400 V
	5Q	(D)	7Q
(A)	$6\varepsilon_0$	(B)	$6\varepsilon_0$
(C)	11Q	(D)	$\frac{Q}{6\epsilon_0}$
of rac	here S_1 of radius r_1 enclos dius r_2 (> r_1) and there are	e no additional char	There is another concentric sphere S
(A)	1:1	(B)	2:1
(C)	1:2	(C)	2:2
differ	ence of 10 V appears be	transferred from or tween the conductor	ne conductor to another, a potentia rs. The capacitance of the conducto
(A)	16×10-8 F	(B)	0.16×10 ⁻⁸ F
(C)	1.6×10 *F	(D)	None of the above
	(C) The f (A) (C) A surflux (A) (C) If 20 poter (A) (C) If chaflux (C) A spli of racelectr (A) (C) Wher differ syster (A)	The force between two free electric flux through S_1 and S_2 and S_3 and S_4 and S_4 are lectric flux of the electric flux through S_1 and S_2 and S_3 and S_4 and S_4 are lectric flux through S_1 and S_2 and S_3 and S_4 and S_4 are lectric flux through S_1 and S_2 and S_3 are lectric flux through S_1 and S_2 and S_3 are lectric flux through S_4 and S_5 and S_6 are lectric flux through S_1 and S_2 and S_3 are lectric flux through S_4 and S_5 and S_6 are lectric flux through S_1 and S_2 and S_3 and S_4 are lectric flux through S_4 and S_5 and S_6 are lectric flux through S_6 and S_6 are lectric flux through S_6 and S_6 and S_6 are lectric flux through S_6 and S_6 are lectrons are difference of S_6 and S_6 are lectrons are difference of S_6 are lectrons are difference of S_6 are lectrons are difference of S_6 and S_6 are lectrons are difference of S_6 are lectrons are difference of S_6 are lectrons are difference of S_6 and S_6 are lectrons are difference of S_6 are lectrons are difference of S_6 and S_6 are lectrons are difference of $S_$	The force between two free electrons spaced 1Å ap (A) $2.3 \times 10^{-8} \text{N}$ (B) (C) $4.6 \times 10^{-8} \text{N}$ (D) A surface element $\overline{\text{ds}} = 5\overline{\text{i}}$ is placed in an electric flux emanating from the surface is (A) 40 units (B) (C) 20 units (D) If 20 J of work has to be done to move an electric potential is 10 V to another point, where potential is (A) 40 V (B) (C) 15 V (D) If charge Q is placed at a distance a/2 above the conflux of the electric field through the square surface (A) $\frac{5Q}{6\varepsilon_0}$ (B) (C) $\frac{11Q}{6\varepsilon_0}$ (B) A sphere S_1 of radius r_1 encloses a total charge Q. To of radius r_2 (> r_1) and there are no additional charge electric flux through S_1 and S_2 is (A) 1:1 (B) (C) 1:2 (C) When 1.0×10^{12} electrons are transferred from or difference of 10 V appears between the conductor system is (A) $16 \times 10^{-8} \text{F}$ (B)

If the radius of a planet is R, and its density is ρ , then escape velocity will be proportional

30.	Wh	at is the color code for a resistor	of resistance S	3 kn with 5% tolerance?
30.	(A)	Green, Orange, Red-gold	(B)	Orange, Green, Red-silver
	(C)	Red, Green, Red-gold	(D)	Orange, yellow, Red-gold
31.	Contract of the Contract of th	noulli's theorem is applicable in t	he case of	
		The compressible liquid in a turb		
		The incompressible liquid in a tur		
		The compressible liquid in a stream		
	(d)	The incompressible liquid in a str	camline flow	
32.	The	current in mA if 2×10 ²⁰ electron	s pass through	a lamp in one minute is
	(A)		(B)	5.33 mA
	(C)	533 mA	(D)	53.3 mA
33.	orbi	Bohr model of hydrogen atom, the it of radius 5.1×10 ⁻¹¹ m at a frequence rent at any point on the orbit of the	cy of 6.8×10**	ving around the nucleus in a circular revolution per second. The equivalent
	(A)	1.088×10 ⁻³ A	(B)	1.088×10 ⁴ A
	(C)	1.088×10 ⁻⁵ A	(D)	1.088×10 ⁴ A
34.	A be	alb of 100W is operated for 6 hou	urs a day. The	units of energy consumed in 7 days
	(A)	4.2 unit	(B)	42 unit
	(C)	420 unit	(D)	none of the above
35.	If the of pa	e mass of proton is approx. 1840 t ath (R _e /R _p) followed in a direction	imes mass of on perpendicular	electron, then the ratio of their radius ar to 'B' will be
	(A)	1:1840	(B)	1840:1
	(C)	2:1840	(D)	1840:2
36.		n distance between two given magnes, where 'k' is	gnetic poles is	halved, force between them become
	(A)	1	(B)	2
	(C)	4	(D)	V4
37.	Maga	netic moment of a current loop be and number of turns is made the	comes k time ree-fold, when	s when diameter of the loop is made re 'k='
	(A)	2	(B)	3
	(C)	6	(D)	12
			The second secon	

30.		The maximum current that can be dra				
	(A)	300 A	(B)	0.3 A		
	(C)	30 A	(D)	60 A		
39.	Whi	ch of the following characteristics of el	ectrons	determine the current in a conductor?		
	(A)	Drift velocity alone	(B)	Thermal velocity alone		
	(C)	Both drift and thermal velocities	(D)	Neither drift nor thermal velocity		
40.	In a	half wave rectifier, the r.m.s. value of	the a.c.	component of the wave is		
	(A)	equal to d.c. value	(B)	more than d.c. value		
	(C)	less than d.c. value	(D)	zero		
41.	C _P a	nd C _v denote the molar specific heats	of a gas	at constant pressure and at constant		
	volu	me respectively. If $\frac{C_p}{C_v} = \gamma$ and $C_p - C_p$	$C_{V} = R. tf$	nen C _v is equal to		
	(A)	$\frac{R}{\gamma-1}$	(B)	$\frac{\gamma-1}{R}$		
	(C)	$\frac{R}{\gamma - 1}$ $\frac{\gamma R}{\gamma - 1}$	(D)	$\sqrt{\frac{R}{\gamma-1}}$		
42.	For a common base amplifier, the values of resistance gain and voltage gain are 3000 and 2800 respectively. The current gain will be					
	(A)	1.1	(B)	0.98		
	(C)	0.93	(D)	0.83		
43.	The	gate for which output is high if at lea	st one in	nput is low		
	(A)	NAND	(B)	NOR		
	(C)	OR	(D)	AND		
44.	The flarge	following four wires are made of the st extension when the same tension i	same n	naterial. Which of them will have the d?		
	(A)	Length=100 cm, diameter=1 mm				
	(B)	Length=200 cm, diameter=2 mm				
	(C)	Length=50 cm, diameter=0.5 mm				
	(D)	Length=300 cm, diameter=3 mm				

45			quid at its	melting point. Freat required for this
		Specific heat	(B)	Latent heat of vaporization
	(A)	External latent heat	(D)	Latent heat of fusion
46.	(C) Let	n _h and n _e be the number of hol conductor. Then	es and co	nduction electrons in an extrinsic
	(A)	$n_b > n_e$	(B)	$n_h = n_e$
	(C)	$n_k < n_k$	(D)	$n_h \neq n_e$
47.	For a	diamagnetic material, which of th	e following	g statement is correct?
	(A)	Magnetic susceptibility > 0	(B)	Magnetic susceptibility < 0
	(C)	Magnetic susceptibility = 0	(D)	Magnetic susceptibility = 1
48.	Two	nuclei have their masses in the ra	itio of 1:3.	The ratio of their nuclear densities
	(A)	13	(B)	1: √3
	(C)	1:1	(D)	3:1
19.	Bohr's	s atomic model explains the		
	(A)	spectrum of hydrogen atom only		
	(B)	spectrum of an atom and/or ions	of one elec	etron only
	(C)	spectrum of hydrogen molecule		
	(D)	none of these		
0.	The ra	adius of the Bohr orbit depends on	which of t	he following?
	(A)	1/n	(B)	n
	(C)	1/n²	(D)	n ²
1.	The st	mallest de-Broglie wavelength amo	ng the Four	particles moving with same velocity
	(A)	N ₂ molecule	(B)	O ₂ molecule
	(C)	Electron	(D)	Proton
2.	Accor	rding to de Broglie's relation if vel-	ocity of par	ticle is infinite, wavelength will be
	(A)	infinite	(B)	small
	(C)	large	(D)	zero

53.	The	The number of ejected photoelectrons from a metal surface increase when							
	(A)	the energy of incident photon	increases						
	(B)	the frequency of incident radia	the frequency of incident radiation increases						
	(C)	the intensity of incident radiat	ion increases						
	(D)	the stopping potential for eject	ted electrons i	ncreases					
54.		minimum energy required for a tocell is called	photoelectron	to escape from a metal surface in a					
	(A)	Stopping voltage	(B)	Planck's constant					
	(C)	Threshold wavelength	(D)	Work function					
55.	Whi	ch of the following is a state fund	ction in therm	odynamics?					
	(A)	Work	(B)	Heat					
	(C)	Enthalpy	(D)	None of the above					
56.	Whi	ch of the following statements ab	out the first l	aw of thermodynamics is true?					
	(A)	It gives the law of conservatio	n of energy						
	(B)	It gives the direction of flow of	f heat						
	(C)	It introduces the concept of en	tropy						
	(D)	It gives the concept of tempera	ature						
57.	Whice load's		of a material	's resistance to deformation under a					
	(A)	Hardness	(B)	Elasticity					
	(C)	Toughness	(D)	Ductility					
58.	What	t is the average velocity of the m	olecules of a	n ideal gas?					
	(A)	Infinity	(B)	Constant					
	(C)	Zero	(D)	Unstable					
59.	A roc of 10	of radius 10 cm and length one 0 kN stretches it along its length	meter is held then the stre	by a clamp at one end and the force ess on the rod is					
	(A)	3.18×10 ⁶ Nm ⁻²	(B)	318×10 ⁶ Nm ⁻²					
	(C)	3.18×10 ⁻⁶ Nm ⁻²	(D)	318×10 ⁻⁶ Nm ⁻²					
0.	Energ	y associated with a one kilogram	m of matter is						
	(A)	9×10 ⁻¹⁶ J	(B)	9×10 ¹⁶ J					
	(C)	90×10 ¹⁶ J	(D)	0.9 ×10 ¹⁶ J					
	Charles and the later of the la								

velocity of light in vacuum h Planck's constant $h (-h/2 \pi)$ e electronic charge $μ_*$ electron magnetic moment $μ_*$ Bohr magneton $μ_*$ nuclear magneton $μ_*$ electron mass m_* proton mass m_* proton mass m_* neutron mass			Chale in vacuum	2.997 924 58 ·108 m/s
h (*h'2 π) $1.054571\cdot 10^{-34}\text{J/s}$ e electronic charge $1.602176\cdot 10^{-19}\text{C}$ μ_s electron magnetic moment $-928.476362\cdot 10^{-26}\text{J/T}$ μ_s Bohr magneton $927.400899\cdot 10^{-26}\text{J/T}$ μ_s nuclear magneton $5.05078317\cdot 10^{-27}\text{J/T}$ m_s electron mass $9.10938188\cdot 10^{-31}\text{kg}$ m_s proton mass $1.67262158\cdot 10^{-27}\text{kg}$ m_s neutron mass $1.67492716\cdot 10^{-27}\text{kg}$ k_s Boltzmann's constant $1.380650\cdot 10^{-23}\text{J/K}$		e		6.626 069 · 10 ³⁴ J/s
h (=h/2π) e electronic charge μ, electron magnetic moment μ, Bohr magneton μ, nuclear magneton π, electron mass π, proton mass π, proton mass λ, Boltzmann's constant 1.602 176 · 10 ⁻¹⁹ C -928.476 362 · 10 ⁻²⁶ J/T 927.400 899 · 10 ⁻²⁶ J/T 927.400 899 · 10 ⁻²⁶ J/T 9.109 381 88 · 10 ⁻²⁷ J/T 9.109 381 88 · 10 ⁻³¹ kg 1.672 621 58 · 10 ⁻²⁷ kg 1.674 927 16 · 10 ⁻²⁷ kg		h		1.054 571 · 10 ⁻³⁴ J/s
## electronic charge ### electron magnetic moment ### Bohr magneton ### nuclear magneton ### proton mass ### proton mass ### neutron mass ### neutron mass ### Bohr magneton ### 1.672 621 58 · 10 ⁻²⁷ kg ### neutron mass ### 1.674 927 16 · 10 ⁻²⁷ kg ### Bohr magneton ### 1.380 650 · 10 ⁻²³ J/K	1/20	h	A SHEWARD	# 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
μ _s Bohr magneton 927.400 899 · 10 ⁻²⁶ J/T μ _s nuclear magneton 5.050 783 17 · 10 ⁻²⁷ J/T m _e electron mass 9.109 381 88 · 10 ⁻³¹ kg m _p proton mass 1.672 621 58 · 10 ⁻²⁷ kg m _N neutron mass 1.674 927 16 · 10 ⁻²⁷ kg k _B Boltzmann's constant 1.380 650 · 10 ⁻²³ J/K	¢		electronic charge	
μ _n Bohr magneton 927.400 899 · 10 · 26 J/T μ _n nuclear magneton 5.050 783 17 · 10 · 27 J/T m _e electron mass 9.109 381 88 · 10 · 31 kg m _p proton mass 1.672 621 58 · 10 · 27 kg m _N neutron mass 1.674 927 16 · 10 · 27 kg k _n Boltzmann's constant 1.380 650 · 10 · 23 J/K	c	u.	electron magnetic moment	-928.476 362 · 10 ⁻²⁶ J/T
μ_N nuclear magneton 5.050 783 17 · 10^{-27} J/T m_e electron mass 9.109 381 88 · 10^{-31} kg m_p proton mass 1.672 621 58 · 10^{-27} kg m_N neutron mass 1.674 927 16 · 10^{-27} kg k_B Boltzmann's constant 1.380 650 · 10^{-23} J/K	В		Bohr magneton	927.400 899 · 10 ²⁶ J/T
m _p proton mass 1.672 621 58 · 10 ⁻²⁷ kg m _N neutron mass 1.674 927 16 · 10 ⁻²⁷ kg k _B Boltzmann's constant 1.380 650 · 10 ⁻²³ J/K	n		nuclear magneton	5.050 783 17 · 10 ⁻²⁷ J/T
m _N neutron mass 1.674 927 16 · 10 ⁻²⁷ kg k _n Boltzmann's constant 1.380 650 · 10 ⁻²³ J/K	el	,	electron mass	9.109 381 88 - 10 ⁻³¹ kg
k _n Boltzmann's constant 1.380 650 · 10 ⁻²³ J/K	þr	,	proton mass	1.672 621 58 · 10 ⁻²⁷ kg
	ne	N	neutron mass	1.674 927 16 · 10 ⁻²⁷ kg
N ₄ Avogadro's constant 6.022 142 · 10 ²³	В		Boltzmann's constant	1.380 650 · 10 ⁻²³ J/K
	Av		Avogadro's constant	6.022 142 · 10 ²³
R molar gas constant $N_A \cdot k_B = 8.314472 \text{ J/mol·K}$	mo		molar gas constant	$N_A \cdot k_B = 8.314472 \text{ J/mol·K}$
F Faraday constant 96 485.3415 C/mol	Far		Faraday constant	

CHEMISTRY-Part A

61.	In th	e modern periodic table, the period	d indicates th	e value of
	(A)	atomic number	(B)	atomic mass
	(C)	principal quantum number	(D)	azimuthal quantum number.
62.	Whic	ch one of the following is an amph	oteric oxide	?
	(A)	Na ₂ O	(B)	SO ₂
	(C)	B ₂ O ₃	(D)	ZnO
63.	Dete	rmine the total number of neutron	s in three iso	topes of hydrogen
	(A)	1	(B)	2
	(C)	3	(D)	4
64.	The	synonym for water gas, when used	in the produ	uction of methanol, is
	(A)	fuel gas	(B)	natural gas
	(C)	laughing gas	(D)	syn gas
65.	Whie	ch of the alkali metal is having lea	st melting p	oint?
	(A)	Na	(B)	K
	(C)	Rb	(D)	Cs
66.	Calc	ulate the number of atoms in 52 m	noles of Ar	
	(A)	52	(B)	3.131×10 ²⁵
	(C)	31.31×10 ²⁵	(D)	1
67.	Calc	ulate the molar mass of H2O in g/	mol	
	(A)	18	(B)	32
	(C)	34	(D)	16
68.	Whi	ch one of the following is tempera	ature indepen	ndent unit of concentration?
	(A)	Molality	(B)	Molarity
	(C)	Normality	(D)	All of the above
69.	Num	ber of unpaired electrons in Mn3*	ion is/are:	
	(A)	1	(B)	2
	(C)	3	(D)	4
	1000111000			

		agnetic moment shown by Cr2 is:		
70		1.00	(B)	3.90
	(A		(D)	2.70
	(C	the structure of diborane the terminal B	—H bon	ads are:
71	(A	- a Jactron bond	(B)	2-centre-2-electron bond
	(C)	3-centre-4-electron bond	(D)	2-centre-4-electron bond
72	W	nich one of the following is the correct	IUPAC r	nomenclature of the compound
-	**	ŅH ₂		
		CH,		
		Ċ ₂ H ₃		
	(A)	4-Ethyl-2-methylaminobenzene		
	(B	1-Methyl-3-ethyl-6-aminobenzene		
	(C)	1-Methyl-2-amino-5-ethylbenzene		
	(D)	4-Ethyl-6-methyl-aniline		
73.	Whi	ch one of the following compounds w	ill not be	e soluble in sodium bicarbonate!
	(A)	Benzene sulphonic acid	(B)	Benzoic acid
	(C)	o-Nitrophenol	(D)	2, 4, 6 - Trinitrophenol
74.		e hydroboration-oxidation reaction of nic compound formed is	propene	with diborane, H ₂ O ₂ and NaOH to
	(A)	(CH ₃) ,COH	(B)	сн,снонсн,
	(C)	СН,СН ₂ ОН	(D)	CH ₃ CH ₂ CH ₂ OH
75.	Basic	rity of H ₃ PO ₄ is:		
	(A)	2	(B)	3
	(C)	4	(D)	1
76.	Which	h of the following has the second hig	hest ele	ctronegativity in the periodic with
	(A)	F	(B)	0
	(C)	N	(D)	c

77	Hig	hest positive oxidation state	of lodine is:	
	(A)	+3	(B)	+5
	(()	+7	(D)	+9
78	Whi	ch one has the highest boil	ing point?	
	(A)	Kr	(B)	Xe
	(C)	He	(D)	Ne
79.	Whi	ch of the following is an in	tensive property?	
	(A)	Mass	(B)	Volume
	(C)	Enthalpy	(D)	Temperature
80.	Mol	ality of 2.5 g of ethanoic ac	aid (CH, COOH) in 7	5 g of benzene
	(A)	0.556 mol kg ⁻¹	(B)	55.6 mol kg ⁻¹
	(C)	0.227 mol kg ¹	(D)	2.5 mol kg ⁻¹
		СНЕ	MISTRY-Par	t B
		(Atten	pt any 30 Quest	ions)
81.	White salts		li metals has the high	est tendency of formation of hydrat
	(A)	Li	(B)	Na
	(C)	К	(D)	Cs
82	Whic	h alkaline earth metal ion	has the highest hyd	ration enthalpy?
	(A)	Be ^{‡+}	(B)	Mg ²⁺
	(C)	Ca2+	(D)	Sr2+
83.	Whic	h of the following is calle	d Caustic Soda ?	
	(A)	NaOH	(B)	NaC1
	(C)	Na ₂ CO ₃ 10H ₂ O	(D)	CaSO ₄
K4	Value	of gas constant R is		
	(A)	0.082 L atm	(B)	0.987 cal mol 1K-1
	(C)	8.3 J mol ¹ K ⁻¹	(D)	83 erg mol ¹ K ¹

85	W	nich one of the following is the wr	ong assumpti	ion of the kinetic theory of gases?
	(A)	All the molecules move in a security.	straight line	between collision and with the same
	(B)	Molecules are separated by gro	eat distances	compared to their sizes.
	(C)	Pressure is the result of the e wall.	lastic collision	on of molecules with the container
	(D)	Momentum and energy always	remain cons	cryed.
86.	Hov	w many electrons an atom may have	ve if the quar	ntum numbers are: n = 3, 1 = 0
	(A)		(B)	10
	(C)	2	(D)	3
87	In th	ne spectrum of electromagnetic ra	diation which	h one have the longest wavelength
	(A)	gamma-rays	(B)	visible rays
	(C)	radio waves	(D)	UV rays
88.	Whi	ch statement is incorrect about the	e Rutherford	Nuclear Model of the atom?
	(A)	most of the α-particles passed		
	(B)	a small fraction of the α-partie		
	(C)			is, were deflected by nearly 180°.
	(D)	The thin foil used in the experi		
9.	Geor	netry of the Molecule CHCl, will		
	(A)	Bent	(B)	T-shape
	(C)	Tetrahedral	(D)	Square-Pyramidal
).	Using	MO theory, predict which of the	following s	pecies has the longest bond length?
	(A)	0,-	(B)	0,2
	(C)	O ₂ ²⁺	(D)	0,
	On tre	ating phenol with chloroform in oduced at the ortho position of the	the presence he benzene r	of sodium hydroxide, a -CHO grouing. This reaction is known as:
	(A)	Reimer-Tiemann Reaction	(B)	Kolbe's Reaction
	(C)	Aldol- Condensation	(D)	Wurtz Reaction
	In Vict	or-Meyer's test, the colour giver	by 1°, 2° a	
		Red, blue, colourless	(B)	
	TERROR III		The T	Colourless, red, blue
((C) 1	Red, blue, violet	(D)	Red, colourless, blue

91

93.	An o	ether is more volatile than alco	phot having the sa	ome molecular formula. This is due
	(A)	alcohols having resonance	structures	
	(B)	inter-molecular hydrogen b		
	(C)	dipolar character of ether	g vinera	
	(D)	inter-molecular hydrogen b	onding in alcohol	
94.	In w	hich of the following ionization agnetic behaviour has change	On processes the	
	(A)	NO → NO'		
	(C)	$N_2 \rightarrow N_2^+$	(D)	$O_2 \rightarrow O_2^*$ $C_2 \rightarrow C_2^*$
95.	The	full form of VSEPR Theory is		
	(A)	Valence shell electron pair	rate theory	
	(B)	Valence shell electron prote		
	(C)	Valence shell electron pair		
	(D)	Valence shell electronegati		theory
96.	Fort	he process to occur under adi	abatic conditions	t, the correct condition is:
	(A)	$\Delta T = 0$	(B)	$\Delta p = 0$
	(C)	q = 0	(D)	W = 0
97.	Whic	th of the following has the hi	ghest bond entha	lpy?
	(A)	C-C	(B)	Si—Si
	(C)	Ge—Ge	(D)	Sn—Sn
98.	An a	queous solution of borax is		
	(A)	neutral	(B)	amphoteric
	(C)	basic	(D)	acidic
99.	The	H of a sample of vinegar is 3	.76. Calculate th	e concentration of hydrogen ion in it;
	(A)	1.7 × 10 ⁻⁴ M	(B)	2.7 × 10 ⁻⁴ M
	(C)	3.76 × 10 ⁻⁴ M	(D)	2.3 × 10 ⁻⁸ M
100.	The r	eaction 3ClO (aq) → ClO,	(aq) + 2Cl ⁻ (aq)	is an example of
	(A)	Oxidation	(B)	Reduction
	(C)	Disproportionation	(D)	Decomposition reaction
			(-)	Decomposition reaction

			THE RESERVE OF THE PARTY OF THE	THE PARTY OF THE P	
101	. The	e pair of compounds having metals	in their high		
	(A)	MnO ₂ , FeCl ₁	(B)	The state of the s	
	100	[Fe(CN) ₆] ³ -, [Co(CN) ₃]		[NiCl ₄] ² , [CoCl ₄]	
102	. EM	F of a cell in terms of the reduction	potential of	fits left and right electrodes	1
	(A)	$E = E_{kit} - E_{right}$	(B)	$\mathbf{E} = \mathbf{E}_{\text{left}} + \mathbf{E}_{\text{right}}$	
	(C)	$E = E_{right} - E_{teft}$	(D)	$\mathbf{E} = -\left(\mathbf{E}_{\text{right}} + \mathbf{E}_{\text{left}}\right)$	
103	. Wh	ich of the following is the chemica	l formula of	sulphurous acid:	
	(A)	H ₂ SO ₃	(B)	H ₂ SO ₄	
	(C)	H ₂ S ₂ O ₇	(D)	H ₂ S ₂ O ₈	
104	. Wh	ich of the following are Lewis acid	s?		
	(A)	AlCl ₃ and SiCl ₄	(B)	PH, and SiCl ₄	
	(C)	BCl ₃ and AlCl ₃	(D)	PH ₃ and BCl ₃	
105.	Wh	ich one of the following types of di	rugs reduces	fever?	
	(A)	Analgesic	(B)	Antipyretic	
	(C)	Antibiotic	(D)	Tranquiliser	
06.	Whi	ich one of the following is the swee	etest artificia	al sweetner?	
	(A)	Aspartame	(B)	Saccharin	
	(C)	Sucralose	(D)	Alitame	
07.	Unit	s of a first-order reaction is:			
	(A)	mol L ⁻¹ s ⁻¹	(B	S ⁻¹	
	(C)	mol ⁻¹ L s ⁻¹	(D)	unitless	
18.	Whie	ch type of 'defect' has the presenc	e of cations	in the interstitial sites?	
	(A)	Vacancy defect	(B)	Frenkel defect	
	(C)	Metal deficiency defect	(D)	Schottky defect	
9.	The I	ydrocarbon which can react with	sodium in l	iquid ammonia is	
	(A)	CH,CH,C≡CCH,CH,			
	(B)	CH,CH,CH,C≡CCH2CH2CH3			
	(1)				
	(C)	CH ₃ CH ₂ C≡CH			
	(D)	CH CH = CHCH.			

110.	The r	caction CH ₂ CH ₂ I + KOH(aq) -	си,си,он	I + KI is classified as
	(A)	electrophilic substitution	(H)	nucleophilic substitution
	(0)	climination	(D)	addition
111	The b	sond order in the molecule NO v	vill be:	
	(A)	2	(B)	1.5
	(C)	2.5	(D)	3
112.	The s	compound with two lone pairs of	electrons on	the central atom is:
	(A)	BrF,	(B)	CIF
	(C)	CH,	(D)	SF ₄
113.	Whic	th one of the following compoun	ds is polar:	
	(A)	0,	(B)	CCI ₄
	(C)	Benzene	(D)	NH ₃
114.	Duri step	ng dehydration of alcohols to al is	kenes by her	nting with conc. H ₂ SO ₄ the initiation
	(A)	formation of carbocation		
	(B)	elimination of water		
	(C)	formation of an ester		
	(D)	protonation of the alcohol mo	lecule	
115.	chlo	e presence of a small amount of rine or bromine to yield a compo- gen. This reaction is known as:	phosphorous ound in which	s, aliphatic carboxylic acids react with ch α – hydrogen has been replaced by
	(A)	Etard reaction		
	(B)	Hell-Volhard-Zelinsky reaction	on	
	(C)	Wolff-Kishner reaction		
	(D)	Rosenmund reaction		
116.	Whi	ch of the following is not an ex-	imple of a h	eterogeneous catalytic reaction?
	(A)	Haber's process		
	(B)	Hydrogenation of vegetable	oils	
	(C)	Combustion of coal		
	(D)	Ostwald's process		

	(A)	81	(B)	C
	(C)	Ge	(D)	Pb
118	e Wh	ich one of the following a	ntibiotic is bacteriost.	atic 1
	(A)	Penicillin		
	(B)	Lrythromycus		
	(C)	Aminoglycosides		
	(D)	Offoxacin		
119	RNA	is different from DNA be	ecause RNA contains	
	(A)	ribose sugar and thymin	ie	
	(B)	ribose sugar and uracil		
	(C)	deoxyribose sugar and t	hymine	
	(D)	deoxyribose sugar and u	racil	
20.	Nylor	threads are made of		
	(A)	Polyester polymer		
	(B)	Polyamide polymer		
	(C)	Polyethylene polymer		
	(D)	Polyvinyl polymer		

MATHEMATICS-Part A

		contraposative of the s	materialit. If p then q" is	
	(A)	If ~p then q	(B)	If p then ~p
	(C)	If q then p	(D)	If -q then -p
122.	The i	range of the function	$f(x) = \frac{2+x}{2-x}, x \neq 2 \text{ is}$	
	(A)	R	(B)	R-{-1}
	(C	R-{1}	(D)	R-{2}
123.	The	function $f: \mathbb{R} \to \mathbb{R}$ gi	iven by $f(x) = - x-1 $	is
	(A)	continuous as well	as differentiable at $x = 1$	
	(B)	not continuous but	differentiable at x = 1	
	(C)	continuous but not	differentiable at $x = 1$	
	(D)	neither continuous	nor differentiable at x =	I management
124.	If A	is a 3×3 matrix such t	hat $ A = 8$, then $ 3A $ eq	uals
	(A)	8	(B)	24
	(C)	72	(D)	216
125.	The	maximum number of	equivalence relations or	the set $A = (1,2,3)$ is
	(A)	1	(B)	2 9 9 9 9 9 9 9 9
	(C)	3	(D)	5
126.	lim[1→2	x] is		
	(A)	2	(B)	3
	(C)	5/2	(D)	5

127	In t	the arithmetic progression $-3, -\frac{1}{2}, 2$.	, the 1	I th term is
	(A)	40	(B)	- 12
	(C)	22	(D)	42
128.	If a	and β are roots of $x^2 + px + q = 0$ the	en the va	due of $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$
	(A)	$\frac{p^2-2q}{q}$	(B)	$\frac{2q-p^2}{q}$
	(C)	$\frac{(p^2+2q)}{q}$	(D)	None of thes
129.	Let	$A = \{x : x \in \mathbb{R}, x < 10\} \text{ and } B = \{x : x \in \mathbb{R} \}$	∈ R, x	> 9}
	The	$A \cap B$ equals to		
	(A)	(9, 10]	(B)	(9, 10)
	(C)	[9, 10)	(D)	[9, 10]
130.	Atp	oint $x = 0$, the function $f(x) = x $ has		
	(A)	Neither minimum nor maximum	(B)	A maxima
	(C)	Point of inflexion	(D)	A minima
131.	The	degree of the differential equation x^2	$\frac{d^2y}{dx^2} = \left[\right]$	$\left(x\frac{dy}{dx}-y\right)^3$ is
	(A)	1	(B)	2
	(C)	3	(D)	6
32.	Whic	h of the following is not a measure o	f central	tendency?
	(A)	Mean	(B)	Median
	(C)	Standard Deviation	(D)	Mode
33.	If q is	the inclination of a line from x-axis,	then its	slope is
	(A)	tan q	(B)	cot q
	(C)	cos q	(D)	$\sin q$

	er feet	tion of the differential equ	uation $xdy - ydx = 0$ re	presents a
134	(A)	parabola	(B)	circle
	100	hyperbola	(D)	straight line
	The I	point which does not lie i	n the half plane $2x + 3$	By - 12 ≤ 0 is
135	(A)	(1, 2)	(B)	
	(C)	(2, 3)		(-3, 2)
36.	a die	is thrown once. Let A be the event that the number	the event that the nur r obtained is less than	mber obtained is greater than 3. Let 5. Then $P(A \cup B)$ is
	(A)	2/5	(B)	3/5
	(C)	0	(D)	1
37.	$\int x^2 e^{-x}$	dx equals		
	(A)	$\frac{1}{3}e^{r^2}+C$	(B)	$\frac{1}{3}e^{x^2}+C$
	(C)	$\frac{1}{2}e^{x^2}+C$	(D)	$\frac{1}{2}e^{x^2}+C$
38.	If 16 /	$P_{r-1}: {}^{15}P_{r-1} = 16:7$, then $P_{r-1}: {}^{15}P_{r-1} = 16:7$	- is	
	(A)	10	(B)	12
	(C)	8	(D)	7
39.	If ā	$\vec{b} = \frac{1}{2} \vec{a} \vec{b} $, then the	angle between \vec{a} and	$1\vec{b}$ is
	(A)	0°	(B)	30°
	(C)	60°	(D)	90°
40.	The r	number of tangents that	can be drawn from (1,2) to the circle
	x2 + y	$y^2 = 5 \text{ is}$		
	(A)	0	(B)	1
	(C)	2	(D)	more than 2

MATHEMATICS-Part B

(Attempt any 30 Questions)

141. The relation R in the set $\{1, 2, 3\}$ given by $R = \{(1, 2), (2, 1), (1, 1)\}$ is

- (A) Symmetric and transitive but not reflexive
- (B) Reflexive and symmetric but not transitive
- (C) Symmetric but neither reflexive nor transitive
- (D) An equivalence relation

142. If
$$g(x) = 1 + x - [x]$$
 and $f(x) = \begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0 \end{cases}$ then for all x .

f(g(x)) is

(A) x

(B) 1

(C) f(x)

(D) g(x)

143. If X and Y are two sets then $X \cap (X \cup Y)^c$ equals

(A) X

(B) Y

(C) 6

(D) None of these

144. The least value of n for which $[(1+i)/(1-i)]^a$ is real is

(A) 1

(B) 2

(C) 3

(D) 4

145. If the roots of the equation $px^2 + qx + 2 = 0$ are reciprocals of each other, then

(A) p=0

(B) p = -2

(C) $p = \pm 2$

(D) p=2

146. If the complex number z = x + iy satisfies the condition |z + 1| = 1, then z lies on

- (A) x-axis
- (B) circle with centre (1,0) and radius 1
- (C) circle with centre (-1, 0) and radius 1
- (D) y-axis

If *P. = 3024 and *C. = 126, then n and r are 147

9,4 (A)

(B) 10, 4

12, 3 (0)

(D) 11, 3

For every natural number k, which of the following statement is true? 145.

 $(mn)^k = m^k n^k$ (A)

(B) $m^k n = mn^k$

 $(m+n)^k = m^k + n^k$ (0)

 $(D) \quad (m-n)^k = m^k - n^k$

The greatest coefficient in the expansion of $(1 + x)^{(n)}$ is 149.

51 (A)

10! (5!×4!) (B)

(C) $\frac{10!}{(5!)^3}$

(D) $\frac{10!}{(4! \times 4!)}$

The principal value of $\cot^4(-\sqrt{3})$ is

(A) $-\frac{\pi}{6}$

(B) $\frac{\pi}{6}$

(C) $\frac{2\pi}{3}$

(D)

151. If $\int_0^a \frac{dx}{1+4x^2} = \frac{\pi}{8}$, then the value of a is

(A) 4

(B)

(C) $\frac{1}{2}$

(D) $\frac{1}{4}$

152. $\int_0^{\pi/3} \tan^2(2x) \, dx$ is equal to

 $(A) \quad \frac{4-\pi}{8}$

(B) $\frac{4+\pi}{8}$

(C) $\frac{4-\pi}{4}$

(D)

153	The interval in which the function f given by f (r) = r ² e	is strictly	increasing is
-----	---	-----------------------	-------------	---------------

154. An urn contains 6 balls of which two are red and four are black. Two balls are draws at random. Probability that they are of different colours is

155. If A and B are two events such that P(A) = 0.2, P(B) = 0.4 and $P(A \cup B) = 0.5$, then the value of P(A/B) is

156. If
$$\begin{vmatrix} 2 & 3 & 2 \\ x & x & x \\ 4 & 9 & 1 \end{vmatrix} + 3 = 0$$
, then the value of x is

157. If $A[a_{ij}]$ is a 2×2 matrix where $a_{ij} = i + j$, then A is equal to

(A)
$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$$

(B)
$$\begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}$$

(C)
$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

(D)
$$\begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}$$

158. Inverse of the matrix
$$\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$$
 is

(A)
$$\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$$

(B)
$$\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$$

(C)
$$\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

(D)
$$\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

159 The existence of the unique solution of the system of equations

$$2x + 3y - z = 6$$

depends on

(A) a only

(B) ß only

(C) Both \alpha and \beta

(D) Neither β nor α

100. If
$$f(x) = \begin{vmatrix} 0 & x-a & x-b \\ x+a & 0 & x-c \\ x+b & x+c & 0 \end{vmatrix}$$
, then

(A) f(a) = 0

(B) f(b) = 0

f(0) = 0

- (D) f(1) = 0
- 161. If P(n) is a statement such that P(3) is true. Assuming P(k) is true implies P(k+1) is true for any integer $k \ge 3$, then P(n) is true for
 - (A) For all $n \in N$

(B) For all $n \ge 3$

(C) For all $n \le 3$

- (D) For all n > 3
- 162. If $f(x) = \begin{cases} ax + 3, x \le 2 \\ a^2x 1, x > 2 \end{cases}$ then the values of a for which f is continuous for all x are
 - (A) 1 and -2

(B) 1 and 2

(C) -1 and 2

(D) -1 and -2

163.
$$\lim_{n \to \infty} \left(\frac{e^{nx} - 1}{e^{nx} + 1} \right)$$
 is

(A) 1

(B) 0

(C) -1

(D) Does not exist

164	The	maximum value of the slope of the cur-	vc y = -	$-x^3 + 3x^2 + 12x - 5$ is
	(A)	15	(B)	12
	(C)	9	(D)	0
165.	The	points of discontinuity of tan x are		
	(A)	$n\pi, n \in I$		
	(B)	$2n\pi, n \in I$		
	(C)	$(2n+1)\frac{\pi}{2}, n \in I$		
	(D)	None of these		
166.	The	condition that $f(x) = ax^3 + bx^2 + cx + d$	has no	extreme value is
	(A)	$b^2 > 3ac$		
	(B)	$b^2 = 4ac$		
	(C)	b ² = 3ac		
	(D)	$b^2 < 3ac$		
67.	If the	first term of a geometric series is 20 a	nd con	amon ratio is 4, then the fifth term is
	(A)	4020	(B)	10240
	(C)	2560	(D)	5120
68.	The s	um of the series $1 + \frac{1}{2} + \frac{1}{4} + \dots$ upto $6^{\frac{1}{2}}$	term is	
	(A)	63	(B)	32
		32		63
	(C)	26	(D)	53
	(-)	53	(0)	26
9.	If y =	$Ae^{5x} + Be^{-5x}$, then $\frac{d^2y}{dx^2}$ is equal to		
	(A)	25y	(B)	5v

(D) 15y

(C) -25y

.70

If x^0 is an integrating factor of the differential equation $\frac{dy}{dx} + Py = Q$, then P can be

(A) 1

(B) $\frac{3}{x}$

(C) 3x

(D) $\frac{2}{x}$

171. The solution of $e^{\frac{dy}{dx}} = x$, when x = 1 and y = 0 is

- (A) $y = x(\log x 1) + 4$
- (B) $y = x(\log x 1) + 3$
- (C) $y = x(\log x + 1) + 2$
- (D) $y = x(\log x 1) + 1$

172. The value of p for which $p(\hat{i} + \hat{j} + \hat{k})$ is a unit vector is

(A) 0

(B) $\frac{1}{\sqrt{3}}$

(C) 1

(D) \sqrt{3}

173. The two lines x = ay + b, z = cy + d and x = a'y + b', z = c'y + d' are perpendicular to each other, if

- (A) $\frac{a}{a'} + \frac{c}{c'} = 1$
- (B) $\frac{a}{a'} + \frac{c}{c'} = -1$
- (C) aa' + cc' = 1
- (D) aa' + cc' = -1

174. ABCD is a rhombus whose diagonal intersect at E, then $\overline{EA} + \overline{EB} + \overline{EC} + \overline{ED}$ equals

(A) 0

(B) <u>AD</u>

(C) 2BC

(D) 2AD

(A) (C) If [a] (A)	2 $4\sqrt{2}$ = 4 and $-3 \le \lambda \le 2$, then $ \lambda $		$2\sqrt{2}$ $3\sqrt{2}$
If [a]			3√2
	= 4 and $-3 \le \lambda \le 2$, then 1λ		
		\hat{a} lies in	
1	[0, 12]		[2, 3]
(C)	[8, 12]	(D)	(-12, 8)
In an	ellipse the distance between stricity is	the foci is 6 and	its minor axis is 8 then its
(A)	4/5	(B)	$\frac{1}{\sqrt{52}}$
(C)	3 5	(D)	$\frac{1}{2}$
f the	mean of the observations: x ,	5, 6, 1, 2 is 4 the	en x is
A)	4	(B)	6
C)	3	(D)	10
oserv	ation is increased by 5, let	indard deviation the new mean a	of the set of observations. If each of the set of observations, if each of the standard deviation be μ_1 and σ_2
A)	$\mu > \mu_1$ and $\sigma = \sigma_1$	(B)	$\mu < \mu_1$ and $\sigma = \sigma_1$
²)	$\mu = \mu_1$ and $\sigma = \sigma_1$	(D)	$\mu_i = \mu_i$ and $\sigma < \sigma_i$
	(A) (C) (f the A) (C) f \(\mu \) abserve hen th	eccentricity is (A) $\frac{4}{5}$ (C) $\frac{3}{5}$ If the mean of the observations: x , (A) 4 (B) 3 If μ and σ are the mean and states the servation is increased by 5, let then the correct option is (A) $\mu > \mu_1$ and $\sigma = \sigma_1$	In an ellipse the distance between the foci is 6 and eccentricity is (A) $\frac{4}{5}$ (B) (C) $\frac{3}{5}$ (D) If the mean of the observations: x , 5, 6, 1, 2 is 4 the A) 4 (B) (C) 3 (D) If μ and σ are the mean and standard deviation bservation is increased by 5, let the new mean and the correct option is (B) (D)

(B.TECH. / B. PHARMACY) 28

If the line 2x - y + a = 0 is a diameter of the circle $x^2 + y^2 + 6x - 6y + 5 = 0$ then a is

(B)

(D)

175

(A)

(C)

5

11

BIOLOGY-Part A

153	Basis	cunit of classification is				
7.	(A)	Species	(B)	Genus		
	(C)	Family	(D)	Phylum		
122.	Aest	ivation is				
	(A)	Arrangement of flowers				
	(B)	Arrangement of sepals and petals in	flower			
	(C)	Arrangement of leaf on stem				
	(D)	All of the above				
123.	Woo	d is				
	(A)	Cambium	(B)	Phloem		
	(C)	Primary xylem	(D)	Secondary xylem		
124.	The	presence of large central vacuote and o	ell wal	I is the feature of		
	(A)	Animal cell	(<u>B</u>)	Plant cell		
	(C)	Bacteria	(D)	Virus		
125.	Poly	peptide chain is made up of following	monon	ner units		
	(A)	Glucose	(B)	Fatty acids		
	(C)	Amino acids	(D)	Nitrogenous bases		
126.	Plasi	nolysis occurs when cell is placed in				
	(A)	Hypotonic solution				
	(B)	Hypertonic solution				
	(C)	Pure water				
	(D)	Isotonic solution				
27.	Which of the following have the capacity to fix atmospheric Nitrogen (N2)?					
	(A)	Angiosperms				
	(B)	Gymnosperms				
	(C)	Bryophytes				
	(D)	Blue Green Algae				

128.	Wh	ich of the following	g is considered as	a compa	tible solute in plants?
	(A)	NaCl		(B)	Proline
	(C)	Both A and B		(D)	None of these
129.	Cole	I treatment required	d to induce flower	ing in pl	ants is called
	(A)	Vernalization			
	(B)	Stratification			
	(C)	Etiolation			
	(D)	None of these			
130.	Whic	h of the following	ion plays major re	ole in the	e stomatal movement?
	(A)	Fe++		(B)	Na+
	(C)	K+		(D)	Zn++
131.	Shive	ring and sweating	are body's way of	regulati	ng
	(A)	Temperature			
	(B)	Water loss			
	(C)	Growth			
	(D)	Metabolism			
132.	What i	s the evolutionary	benefit of light sl	kin in the	e northern latitude?
((A)	Skin cancer protec	ction		
- (B)	Folate protection			
(C)	Easier vitamin D p	production		
(1		No evolutionary re			
33. T		micals that can ca	EL 1997 EL 1997 EN 1997 EL 199	lled	
(A		Veurotoxins			C:
(C		'ytotoxins		(B)	Carcinogens
	-			(D)	Poisons
		id is an example of	of which type of	joint?	
(A) C	artilaginous		(B)	Synovial
(C)) Fi	brous		(D)	None of these

135.	Which of the following will have the maximum heart beat rate?							
	(A)	Human	(B)	Horse				
	(C)	Elephant	(D)	Mouse				
136.	The	nature of nerve impulse conduction	is					
	(A)	Mechanical	(B)	Thermal				
	(C)	Electrochemical	(D)	Chemical				
137.	Repl	Replicate the following strand of DNA: AATCATGGA						
	(A)	UUAGUACCU	(B)	TTAGTACCT				
	(C)	AATCATGGA	(D)	GGATAUCUA				
138.	Follo	owing is the example of sex-linked	inheritance					
	(A)	Colour blindness	(B)	Haemophilia				
	(C)	Both A and B	(D)	None of the above				
139.	Gree	at Himalayan national park and Pin	Valley Nat	ional park are situated at				
	(A)	Kullu and Spitti	(B)	Lahaul and Shimla				
	(C)	Mandi and Kangra	(D)	Chamba and Kullu				
140.	Pyramid of energy of an ecosystem will always be							
	(A)	Upright	(B)	Inverted				
	(C)	Spindle shaped	(D)	None of these				
	BIOLOGY-Part B							
		(Attempt any	30 Que	stions)				
141.	In fiv	e kingdom classification system,	Monera in	cludes				
	(A)	All unicellular eukaryotes						
	(B)	All prokaryotes						
	(C)	Both A and B						
	m	Mark Company of the C						

142	. Wh	ich of the following is called as the an	ophibian o	of the plant kingdom?
	(A)	Algae	(B)	Bryophytes
	(C)	Pteridophytes	(D)	Gymnosperms
143.	Liel	nens are		
	(A)	Bryophytes		
	(B)	Algae		
	(C)	Fungi		
	(D)	An association between algae and	fungi	
144.	Whi	ch of the following is an example of	"living fo	ssil"?
	(A)	Gnetum	(B)	Ginkgo
	(C)	Pinus	(D)	Taxus
145.	The	largest phylum in the kingdom Anim	alia is	
	(A)	Mollusca	(B)	Annelida
	(C)	Nematoda	(D)	Arthropoda
146	Can	al system is present in phylum		
	(A)	Porifera	(B)	Echinodermata
	(C)	Protozoa	(D)	Cnidaria
147.	Whic	th of the following is a vertebrate?		
	(A)	Cuttle fish	(B)	Cray fish
	(C)	Trout fish	(D)	Silver fish
48.	The c	edible part of Litchi is		
	(A)	Ovary	(B)	Thalamus
	(2)	Aril	(D)	Cotyledons
49	1-2	edons are main food storing organ		
		Wheat		
	(A)		(B)	Maize
	(C)	Bean	(D)	Barley

		mosmotic hypothesis of ATP synthesis	s was giv	en by				
150	*	Robert Hill	(H)	Calvin				
	(A)	Peter Mitchell	(D)	Levitt				
	2000	is an example of						
151	(A)	Plant Growth regulator						
	(B)	Anti-transpirant						
	(C)	Plant stress hormone						
	(D)	All of the above						
	200	suc culture *Callus* refers to						
152	(A)	Mass of undifferentiated cells						
	(B)	Differentiated cells						
	(C)	Root formation						
	(D)	Shoot formation						
153.	Crossing over occurs between							
122		(A) Non-sister chromatids during leptotene						
	(B)	to chromatids during Pachytene						
	(c)	Sister chromatids during Pachyten	ie					
	(D)	Sister chromatids during Zygotene						
	The	wo strands of DNA double helix are	attache	d to each other through				
154.		Covalent bond	(B)	Hydrogen bond				
	(A)		(D)	Disulfide bond				
	(C)	Ionic bond	Administra					
155.	The c	The concept of Operon model was given by						
	(A)	Hershey and Chase						
	(B)	Jacob and Monod						
	(c)	Ruben and Kamen						
	(D)	Lwaff and Went						

156. What part of a nucleotide accounts	for the genetic	e variation between individuals?
(A) Nitrogenous base		
(B) Deoxyribose		
(C) Phosphate		
(D) All of above		
157. Nobel Prize for the discovery of do	uble helix of I	NA molecule was awarded to
(A) Watson and Crick		
(B) Watson and Wilkins		
(C) Wilkins and Crick		
(B) Watson, Crick and Wilkins		
158. Which of the following is a character	eristics of smo	oth muscle cells?
(A) Voluntary	(B)	Striated
(C) Non-striated	(D)	Multinucleate
159. Which of the following is common	to aerobic and	anaerobic respiration?
(A) Glycolysis		
(B) Krebs cycle		
(C) Electron transport chain		
(D) All of above		
160; Starch is digested by		
(A) Protease	(B)	Amylase
(C) Lipase	(D)	Catalase
Vermiform appendix is a part of		
(A) Liver	(B)	Stomach
(C) Intestine	(D)	Rectum
162. One molecule of haemoglobin carries	how many m	olecules of oxygen?
(A) Two	(B)	Four
	(D)	Eight
(C) Six	. (D)	Light

	-	nd product of ormanne cycle is					
103		Ammonia	(B)	Urea			
	(A)	Uric acid	(D)	Ethanol			
	(C)	regulation is the function of					
164.	(A)	Oxytocin	(V)	Prolactin			
ing wit	(0)	Insulin	(D)	Vasopressin			
	Which	h of the following organelle he	elps the sperm t	o penetrate the ovum?			
To.	(A)	Acrosome	(B)	Zona pellucida			
	(C)	Glyoxysome	(D)	Ribosome			
102	Whic	h of the following process is re	elated to ATP s	ynthesis?			
166.	(A)	Substrate level phosphorylat	ion				
	(B)	Photophosphorylation					
	(C)	Oxidativephosphorylation					
	(D)	All of these					
167.		opterophily refers to					
200	(A)	Insect pollination	(B)	Bat pollination			
	(C)	Water pollination	(D)	Air pollination			
150	Whic	h of the following is related to	o silk production	on?			
P	(A)	Silviculture	(B)	Tissue culture			
	(C)	Sericulture	(D)	None of these			
169.	The	theory of 'Natural selection' w	as given by?				
•	(A)	Lamarck	(B)	Darwin			
	(C)	Wallace	(D)	Spencer			
170.	Hom	ologous organs are					
	(A)						
	(B)	Dissimilar in origin but sin		n			
	(C)	Similar in structure and fur					
	(D)	Discimilar in origin and fur					
		The state of the s					

171 1	Individual with Turner's syndrome is					
(A)	Normal male				
(B)	Normal female				
(C)	A male with rudimentary testis and u	nderde	veloped penis		
0	D)	A female with rudimentary ovaries a	nd unde	erdeveloped breasts		
172. V	Vhic	h of the following study provides evid	ences ir	favour of biological evolution?		
(4	A)	Archaeology	(B)	Paleontology		
((C)	Phycology	(D)	Mycology		
173. N	on-s	ense codon codes for				
(4	1)	Proline	(B)	Lysine		
(0)	Tryptophan	(D)	None of these		
174. TI	ne st	rength of linkage depends on the				
(A	()	Distance between linked genes				
(B)	Length of chromosomes				
(C)	Size of genes				
(D)	None of these				
175. Mi	nam	ata disease is caused by				
(A)) 1	Lead	(B)	Silver		
(C)	1	Mercury	(D)	Nitrogen		
76. Het	егос	yst is a specialized cell present in				
(A)	1	lostoc	(B)	Azolla		
(C)	(yeus	(D)	Pinus		
2. Che	motl	herapy and radiation therapy are ger	nerally	used to treat		
(A)	A	IDS	(B)	Cancer		
(C)	Н	aemophilia	(D)	Tuberculosis		

178	Whic	h of the following gases c	ause Greenhouse e	ffect?	
Y	(A)	Carbon dioxide			
	(B)	Methane			
	(C)	Nitrous oxide			
	(D)	All of these			
1000	Gold defic	en rice is a genetically mo iency of dietary	dified rice variety o	developed to overcome the	
	(A)	Protein	(B)	Iron	
	(C)	Vitamin A	(D)	All of the above	
180.	Which of the following indicates the higher level of population threat status of a species?				
	(A)	Critically Endangered			
	(B)	Endangered			
	(C)	Vulnerable			
	(D)	Low risk near threatened	d		